

(Defended January 17, 2008)

© 2008

Brian M Zid

All Rights Reserved

Acknowledgements

I would like to thank Seymour Benzer for taking me into his lab, giving me guidance and at the same time, freedom in pursuing my research. I especially value the insight I have gained from him on pursuing novel ideas in science and hope that this knowledge will guide me in my future research endeavors. I would like to thank Pankaj Kapahi for always being there to bounce ideas off of and argue with, and for bringing excitement to the science, even if the experiments weren't always going as expected. I would like to thank Eimear Kenny for listening and helping me scientifically as well as non-scientifically during my time at Caltech. I would like to thank my committee members, Ray Deshaies, Alex Varshavsky, Judy Campbell and, especially, Paul Sternberg for always having their doors open to answer my questions and to have scientific discussions. I would also like to thank members of the Benzer and Kapahi Labs as well as other members of the Caltech community for all their help.

Abstract

Aging is characterized by the declining ability of an organism to maintain homeostasis, which eventually leads to death. Dietary restriction (DR), the reduction of nutrients without malnutrition, extends lifespan in various organisms, yet its molecular underpinnings are poorly understood. We show that in *Drosophila*, DR upregulates the translational repressor 4EBP, the eukaryotic translation initiation factor 4E binding protein, and that this upregulation is necessary for the full lifespan extension upon DR and sufficient to extend lifespan on a nutrient rich diet. Investigation of the genome-wide translational changes upon DR using translation state array analysis (TSAA) found that translationally downregulated genes tend to have extensive 5' untranslated regions (UTR) secondary structures, while those that are upregulated have weakly structured 5'UTRs. Among the translationally upregulated genes, mitochondrial ribosomal proteins and electron transport chain components were overrepresented. Mitochondrial genes were found to have weakly structured 5'UTRs in *Drosophila*, and this was conserved in Humans. The 5'UTRs of mitochondrial genes were found to be sufficient to confer preferential translation during times of high 4EBP activity in a cap-independent manner to reporter constructs. Upregulation of mitochondrial function was verified and found to be *d4EBP* dependent, implicating a novel mechanism for regulating mitochondrial function upon DR. These results implicate mRNA translation initiation in modulating lifespan and mitochondrial function upon DR.

Contents

Acknowledgements	iii
Abstract	iv
Background	1
1 Aging	1
1.1 Introduction	1
1.2 Dietary Restriction	1
1.3 Genetics of Aging	2
1.31 Insulin Pathway	2
1.32 TOR Pathway	3
1.4 Genetics of Dietary Restriction	5
1.41 Sir2	5
1.42 TOR Pathway	6
1.43 Insulin Pathway	6
1.5 Mitochondrial Respiration	7
2 Translation	8
2.1 Translation Initiation	8
2.11 Cap Dependent Translation Initiation	8
2.12 Internal Ribosome Entry	10
2.13 Cancer	10
2.2 Lifespan	11
Results	11

3 Introduction	11
4 4EBP and Dietary Restriction	13
4.1 Protein Levels	14
4.2 Lifespan	14
4.21 4EBP Null	14
4.22 4EBP Overexpression	20
4.3 Metabolic Changes	22
4.4 Stress Resistance	24
5 Genome Wide Translational Changes	25
5.1 Translation State Array Analysis	25
5.2 5'UTR Analysis	27
5.3 Mitochondrial Measurements	30
6 5'UTR Analysis	32
6.1 Conserved Gene Ontology Categories	32
6.2 In vivo Analysis of 5'UTR Function	36
6.21 Monocistronic Reporter	37
6.22 Bicistronic Reporter	39
Concluding Remarks	40
7 Summary of Results	40
8 Discusion.....	42
8.1 IRES Translation	42
8.2 Growth and Lifespan	44
8.21 Multicellular Organisms	44

8.22 Yeast Growth Changes	45
8.23 Mitochondrial Respiration.....	46
Methods	49
Supplemental Tables	57
References	60